















- [30] C.-H. J. Tzou, N. M. Artner, I. Pona, A. Hold, E. Placheta, W. G. Kropatsch, und M. Frey, „Comparison of three-dimensional surface-imaging systems“, *J. Plast. Reconstr. Aesthet. Surg.*, Bd. 67, Nr. 4, S. 489–497, Apr. 2014.
- [31] T. Beeler, „Passive Spatiotemporal Geometry Reconstruction of Human Faces at High Fidelity“, *IEEE Comput. Graph. Appl.*, Bd. 35, Nr. 3, S. 82–90, Mai 2015.
- [32] N. D'Apuzzo, „Modeling human faces with multi-image photogrammetry“, in *Electronic Imaging 2002*, 2002, S. 191–197.
- [33] M. Abreu de Souza, S. Robson, und J. C. Hebden, „A Photogrammetric Technique for Acquiring Accurate Head Surfaces of Newborn Infants for Optical Tomography Under Clinical Conditions“, *Photogramm. Rec.*, Bd. 27, Nr. 139, S. 253–271, Sep. 2012.
- [34] I. Detchev, M. Mazaheri, S. Rondeel, und A. Habib, „Calibration of multi-camera photogrammetric systems“, *ISPRS - Int. Arch. Photogramm. Remote Sens. Spat. Inf. Sci.*, Bd. XL-1, S. 101–108, Nov. 2014.
- [35] M. Pesce, L. M. Galantucci, G. Percoco, und F. Lavecchia, „A Low-cost Multi Camera 3D Scanning System for Quality Measurement of Non-static Subjects“, *Procedia CIRP*, Bd. 28, S. 88–93, 2015.
- [36] M. M. de Sainte Croix, D. Gauld, A. H. Forgie, und R. Lowe, „Three-dimensional imaging of human cutaneous forearm bite marks in human volunteers over a 4 day period“, *J. Forensic Leg. Med.*, Bd. 40, S. 34–39, Mai 2016.
- [37] A. Leipner, R. Baumeister, M. J. Thali, M. Braun, E. Dobler, und L. C. Ebert, „Multi-camera system for 3D forensic documentation“, *Forensic Sci. Int.*, Bd. 261, S. 123–128, Apr. 2016.
- [38] F. Remondino und S. El-Hakim, „Image-based 3D Modelling: A Review“, *Photogramm. Rec.*, Bd. 21, Nr. 115, S. 269–291, Sep. 2006.
- [39] F. Remondino, S. F. El-Hakim, A. Gruen, und L. Zhang, „Turning images into 3-D models“, *IEEE Signal Process. Mag.*, Bd. 25, Nr. 4, S. 55–65, Juli 2008.